

A LIST OF PUBLICATIONS SENT FROM JAPANESE AQUACULTURE PANEL, UJNR, TO
U. S. PANEL AS AN INFORMATION EXCHANGE FOR 1982.

1. Bulletin of National Research Institute of Aquaculture, no. 1 (1980).
2. Bulletin of National Research Institute of Aquaculture, no. 2 (1981).
3. Bulletin of the Seikai Regional Fisheries Research Laboratory, no. 51 (1978).
4. Bulletin of the Seikai Regional Fisheries Research Laboratory, no. 52 (1979).
5. Bulletin of the Seikai Regional Fisheries Research Laboratory, no. 53 (1979).
6. Bulletin of the Seikai Regional Fisheries Research Laboratory, no. 54 (1980).
7. Bulletin of the Seikai Regional Fisheries Research Laboratory, no. 55 (1980).
8. Bulletin of the Seikai Regional Fisheries Research Laboratory, no. 56 (1981).
9. Tanaka, K. and T. Tanaka. 1980. On the age and the growth of Japanese abalone Nordotis discus, in the coast of Chiba Prefecture. , Bull. Japan Sea Region. Fish. Res. Lab., 31 : 115-127.
10. Tanaka, K. 1980. On the mass survivors of Japanese scallop, Pecten (Notovola) albicans, in Tateyama Bay., Bull. Japan Sea Region. Fish. Lab., 31 : 165-168.
11. Nakanishi, T. and M. Naryu. 1981. Some aspects of large-scale rearing of larvae and post-larvae of the king crab (Paralithodes camtschatica), Bull. Japan Sea Region. Fish Res. Lab., 32: 39-47.
12. Nakanishi, T. 1981. The effect of temperature on growth, survival and oxygen consumption of larvae and post-larvae of Paralithodes brevipes (Decapoda : Anomura), Bull. Japan Sea Region. Fish. Res. Lab., 32: 49-56.
13. Akamine, T. 1981. A handy method for selecting and counting bivalve larvae at planktonic stage., Bull. Japan Sea Region. Fish. Res. Lab., 32: 77-81.
14. Morioka, Y. 1981. Zooplankton production in the Toyama Bay in March - May, 1978. , Bull. Japan Sea Region. Fish. Res. Lab., 32: 57-64.
15. Koshiishi, Y. 1981. On the feed conversion efficiency of juvenile plaice, Paralichthys olivaceus., Bull. Japan Sea Region. Fish. Res. Lab., 32: 73-75.
16. Uki, N. 1981. Food value of marine algae of order Laminariales for growth of the abalone, Haliotis discus HANNAI., Bull. Tohoku Region. Fish. Res. Lab., 42:19-29.
17. Seki, T. and H. Kan-no. 1981. Induced settlement of the Japanese abalone, Haliotis discus, veliger by the mucous trails of the juvenile and adult abalones., Bull. Tohoku Region. Fish. Res. Lab., 43: 29-36.

18. Uki, N. and S. Kikuchi. 1981. Technical study on artificial spawning of abalone, genus Haliotis discus VII. Comparative examinations of rearing apparatus for conditioning adult of abalone., Bull. Tohoku Region. Fish. Res. Lab., 43: 47-51.
19. Uki, N. 1981. Feeding behavior of experimental populations of the abalone, Haliotis discus HANNAI., Bull. Tohoku Region. Fish. Res. Lab., 43: 53-58.
20. Uki, N., J. F. Grant and S. Kikuchi. 1981. Juvenile growth of the abalone, Haliotis discus HANNAI, fed certain benthic micro algae related to temperature., Bull. Tohoku Region. Fish. Res. Lab., 43: 59-64.
21. Fujii, T. 1981. The study for periodic behaviour of bivalve - II. Shell movement of ark shell and Pacific oyster., Bull. Tohoku Region. Fish. Res. Lab., 43:65-69.
22. Yamamoto, M., et al. 1981. Water flow characteristics in Hamada Fishing Port., Fisheries Engineering, 18: 31-38.
23. Shimozake, Y. and Y. Osawa. 1981. On the experimental machine devised for determination in breaking stress of webbings and the result tested., Bull. Natl. Res. Inst. Fish. Engineer., 2: 105-112.
24. Hideshima, Y. 1981. Observation of internal tide in Nomi Bay., Fisheries Engineering, 18: 39-48.
25. Toda, S. and M. Nakamura. 1981. Experimental study on wave-induced circulation related to offshore structure., Bull. Natl. Res. Inst. Fish. Engineer., 2: 1-11.
26. Sasaki, M. and S. Sato. 1969. Composition of medium and cultural temperature of Pythium sp., a pathogenic fungus, of the "Akagusare" disease of cultivated Porphyra., Bull. Tohoku Region. Fish. Res. Lab., 29: 125-132.
27. Sasaki, M. and Y. Sakurai. 1972. Comparative observation on the growth among the five strains in Pythium porphyrae under the same cultural condition., Bull. Tohoku Region. Fish. Res. Lab., 32: 83-87.
28. Taniguchi, K., H. Kito and K. Akiyama. 1981. Morphological variation of Undaria pinnatifidae (HARVEY) SURINGAR - I. On the difference of growth and morphological characteristics of two types at Matsushima Bay, Japan., Bull. Tohoku Region. Fish. Res. Lab., 42: 1-9.
29. Kito, H., K. Taniguchi and K. Akiyama. 1981. Morphological variation of Undaria pinnatifida (HARVEY) SURINGAR - II. Comparison of the thallus morphology of cultured F₁ plants originated from parental types of two different morphologies., Bull. Tohoku Region. Fish. Res. Lab., 42:11-18.
30. Akashige, S. et al. 1981. Effects of γ -aminobutyric acid and certain neurotransmitters on the settlement and the metamorphosis of the larvae of Haliotis discus HANNAI INO (Gastropoda)., Bull. Tohoku Region. Fish. Res. Lab., 43: 37-45.

31. Takahashi, M, T. Ichitani and M. Sasaki. 1977. Pythium porphyrae TAKAHASHI ET SASAKI, sp. nov. causing red rot of marine red algae Porphyra spp., Transact. Mycol. Soc. Japan, 18: 279-285.
32. Kasahara, S. and T. Akiyama. 1976. Notes on the dormancy in the adults of Tigriopus japonicus., J. Fac. Fish. Anim. Husb., Hiroshima Univ., 15: 57-65.
33. Akiyama, T. and T. Nose. 1980. Changes in body weight, condition factor and body composition of fingerling chum salmon with various sizes during starvation., Bull. Natl. Res. Inst. Aquaculture, 1: 71-78.
34. Akiyama, T., I. Yagisawa and T. Nose. 1981. Optimum levels of dietary crude protein and fat for fingerling chum salmon., Bull. Natl. Res. Inst. Aquaculture, 2: 35-42.
35. Akiyama, T., A. A. Kajukulo and S. Olsen. 1977. Seasonal variations of plankton and physicochemical condition in Mwanza Gulf, Lake Victoria., Bull. Freshwater Fish. Res. Lab., 27: 49-61.
36. Shimma, Y. et al. 1977. On the lipids of adult carps raised on fish meal and SCP feeds, and hatchabilities of their eggs., Bull. Freshwater Fish. Res. Lab., 27: 35-48.
37. Shimma, Y. and H. Shimma. 1980. Influence of the commercial feed oil supplement to a fish meal ration on body and fatty acid composition of Ayu, Plecoglossus altivelis., Bull. Tokai Region. Fish. Res. Lab., 103: 83-91.
38. Shimma, Y., H. Shimma and K. Ikeda. 1980. Effects of supplemental oils to single cell protein feeds on the growth and fatty acid composition of Ayu, Plecoglossus altivelis., Bull. Natl. Res. Inst. Aquaculture, 1: 47-60.
39. Shimma, Y., K. Ikeda and T. Maruyama. 1980. Fatty acid composition of the trunk flesh and eggs of Ayu, Plecoglossus altivelis, fed single protein feeds., Bull. Natl. Res. Inst. Aquaculture, 1: 61-69.
40. Shimma, Y., K. Ikeda and H. Shimma. 1981. Plasma constituents of matured rainbow trout raised with fish meal and SCP combined feeds., Bull. Japan. Soc. Sci. Fish., 47: 371-376.
41. Murai, T. et al. 1980. Protein, fat and carbohydrate sources of practical diet for fingerling chum salmon, Oncorhynchus keta., Bull. Natl. Res. Inst. Aquaculture, 1: 79-86.
42. Murai, T., Y. Hirasawa and T. Nose. 1981. Optimum supplemental level of vitamin premix in a practical diet for fingerling chum salmon, Oncorhynchus keta., Bull. Natl. Res. Inst. Aquaculture, 2: 43-47.
43. Tanaka, N., M. Sugiyama and K. Ohwada. 1981. Bacterial population and their growth rates in fish rearing tank with running water., Bull. Natl. Res. Inst. Aquaculture, 2: 65-71.
44. Murai, T., T. Akiyama and T. Nose. 1981. Use of crystalline amino acid coated with casein in diets for carp., Bull. Japan. Soc. Sci. Fish., 47: 523-527.

45. Murai, T., H. Ogata and T. Nose. 1982. Methinine coated with various materials supplemented to soybean meal diet for fingerling carp Cyprinus carpio and channel catfish Ictalurus punctatus., Bull. Japan. Soc. Sci. Fish., 48:85-88.
46. Suga, S., K. Wada and M. Ogawa. 1981. Fluoride concentration in teeth of tetraodontiform fishes and its phylogenetic significance., Japan. J. Ichthy., 28: 304-312.
47. Arai, S. 1981. A purified test diet for coho salmon, Oncorhynchus kisutch, fry. Bull. Japan. Soc. Sci. Fish., 47: 547-550.
48. Takeuchi, T. et al. 1980. Requirement of eel Anguilla japonica for essential fatty acids., Bull. Japan. Soc. Sci. Fish., 46: 345-353.
49. Ishida, R. and Y. Shimma. 1979. Effects of single cell protein feeds on the growth and fatty acid composition of ayu, Plecoglossus altivelis., Bull. Tokai Region. Fish. Res. Lab., 100: 161-167.
50. Tanaka, Y. 1978. Culture experiments on growth of Batillus cornutus (gastropoda) and Anthocidaris crassispina (sea urchin) in seawater of high temperature levels., in "Results of Environmental Assessment on the Effects of Heated Discharge Water on Organisms in Coastal Waters", Fisheries Agency, 22-29.
51. Tanaka, Y. 1949. Injuring mechanisms of oyster drill (Purpura clavigera) to young oyster (Ostrea gigas)., Bull. Japan. Soc. Sci. Fish., 15: 447-457.
52. Tanaka, Y. 1970. Effects of sewage sludge upon the growth and mortality of some bivalve mollusks., Bull. Tokai Region. Fish. Res. Lab., 63: 65-74.
53. Tanaka, Y. 1969. Studies on reducing mortality of larvae and juveniles in the course of the mass-production of seed abalone - I. Satisfactory result with streptomyci to reduce intensive mortality., Bull. Tokai Region. Fish. Res. Lab., 58: 155-161.
54. Tanaka, Y., S. Inoha and K. Kakazu. 1970. Studies on seed production of black-lip pearl-oyster, Pinctada margaritifera, in Okinawa - I. Spawning-induction by thermal-stimulation., Bull. Tokai Region. Fish. Res. Lab., 63: 75-78.
55. Tanaka, Y., S. Inoha and K. Kakazu. 1970. Studies on seed production of black-lip pearl-oyster, Pinctada margaritifera, in Okinawa - II. Food micro-organisms available for the larvae., Bull. Tokai Region. Fish. Res. Lab., 63: 79-85.
56. Tanaka, Y., S. Inoha and K. Kakazu. 1970. Studies on seed production of black-lip pearl-oyster, Pinctada margaritifera, in Okinawa - III. Culture experiment of Monochrysis lutheri at high water temperature levels., Bull. Tokai Reg. Fish. Res. Lab., 63: 87-90.
57. Tanaka, Y., S. Inoha and K. Kakazu. 1970. Studies on seed production of black-lip pearl-oyster, Pinctada margaritifera, in Okinawa - IV. Resistibility of the larvae to centrifugally separated water from Monochrysis culture and to hypotonic sea water., Bull. Tokai Region. Fish. Res. Lab., 63: 91-95.

58. Tanaka, Y., S. Inoha and K. Kakazu. 1970. Studies on seed production of black-lip pearl-oyster, Pinctada margaritifera, in Okinawa - V. Rearing of the larvae., Bull. Tokai Region. Fish. Res. Lab., 63: 97-106.
59. Tanaka, Y. 1970. Studies on molluscan larvae - II. Pinctada margaritifera., Venus, 29: 117-122.
60. Tanaka, Y. 1954. Spawning season of important bivalves in Ariake Bay -I. Anadara subcrenata (LISCHKE)., Bull. Japan. Soc. Sci. Fish., 19: 1157-1160.
61. Tanaka, Y. 1954. Spawning season of important bivalves in Ariake Bay - II. Ostrea rivularis GOULD and O. gigas THUNBERG., Bull. Japan. Soc. Sci. Fish., 19: 1161-1164.
62. Tanaka, Y. 1954. Spawning season of important bivalve in Ariake Bay - III. Venerupis semidecussata (REEVE)., Bull. Japan. Soc. Sci. Fish., 19: 1165-1167.
63. Tanaka, Y. 1959. Spawning season of important bivalves in Ariake Bay -IV. Anadara ursus KURODA MS., Report of the Investigations on the Ariake Sea. Seikai Region. Fish. Res. Lab., 5: 1-3.
64. Tanaka, Y. 1981. Identification of bivalve larvae - (13), Aquabiology, 3: 200-203.
65. Tanaka, Y. 1981. Identification of bivalve larvae - (14), Aquabiology, 3: 310-313.
66. Tanaka, Y. 1981. Identification of bivalve larvae - (15), Aquabiology, 3: 387-390.
67. Tanaka, Y. 1982. Identification of bivalve larvae - (16), Aquabiology, 4: 23-26.
68. Tanaka, Y. 1969. Studies on propagation of a hard clam, Meretrix lamarckii - I. Artificial breeding., Bull. Tokai Region. Fish. Res. Lab., 58: 163-168.
69. Tanaka, Y. 1969. Studies on propagation of a hard clam, Meretrix lamarckii - II. Resistibility to hypotonic sea water., Bull. Tokai Region. Fish. Lab., 58: 169-172.
70. Tanaka, Y. 1971. Studies on propagation of a hard clam, Meretrix lamarckii - III. Identification of the larva and spat., Bull. Tokai Region. Fish. Res. Lab., 64: 101-109.
71. Yano, I. 1981. Scanning electron microscopic study on the comb-like teeth of Plecoglossus altivelis TEMMINCK et SCHLEGEL., Bull. Japan. Soc. Sci. Fish., 47: 437.
72. Shimma, H. and A. Tsujigado. 1981. Some biochemical qualities of bred scorpenoid fish, Sebastiscus marmoratus, and activities of their larvae., Bull. Natl. Res. Inst. Aquaculture, 2: 11-20.

73. Tanaka, Y. and M. Kumeta. 1981. Successful artificial breeding of the silver-lip pearl oyster, Pinctada maxima (JAMESON)., Bull. Natl. Res. Inst. Aquaculture, 2: 21-28.
74. Yano, I. 1981. Effects of sea water of reduced salinity on positive phototaxis and negative geotaxis of abalone larvae, Haliotis gigantea., Bull. Natl. Res. Inst. Aquaculture, 2: 29-34.
75. Yano, I. and Y. Ogawa. 1981. Swimming speeds of larvae and juvenile of red sea bream, Chrysophrys major., Bull. Natl. Res. Inst. Aquaculture, 2: 49-54.
76. Kiso, K. 1980. On the feeding habit of 0-group red sea bream (Pagrus major) in Shijiki Bay, Hirado Island - I. Sequential changes of the diet with growth and its annual variation., Bull. Seikai Region. Fish. Res. Lab., 54: 291-306.
77. Kiso, K. and Y. Matsumiya. 1980. Tagging experiments for estimating abundance of 0-group red sea bream, Pagrus major, in Shijiki Bay, Hirado Island., Bull. Seikai Region. Fish. Res. Lab., 54: 307-314.
78. Kiso, K. 1981. On the feeding habit of 0-group crimson sea bream, Evynnis japonica, in Shijiki Bay, Hirado Island., Bull. Seikai Region. Fish. Res. Lab., 56: 15-26.