

A List of Publication from Japanese Aquaculture Panel, UJNR, to U.S. Panel as an Information Exchange for 1988

- 1)Fukusho,K., Nanba,H., Yamamoto,T.,Yamasaki,M., Lee,M.T., Seikai,T. and Watanabe,T.
1987. Reduction of albinism in juvenile flounder Paralichthys olivaceus hatchery-reared on fertilized eggs of red bream Pagrus major, and its critical stage for the effective feeding. Bull. Natl. Res.Inst. Aquaculture 12:1-7
- 2)Wada,K.T.
1987. Selective breeding and intraspecific hybridization - molluscs
In: Selection, Hybridization and Genetic Engineering in Aquaculture, Bordeaux 27-30 May, 1986. Vol. I. Berlin 1987.
- 3)Nakamura,H.K., Machii,A. Wada,K.T. Awaji,M. and Townsley S.J.
1988. A check list decapod chromosomes (Crustacea). Bull. Natl. Res. Inst. Aquaculture 13:1-9.
- 4)Komaru,A., Uchiyama,Y., Ieyama,H. and Wada,K.T.
1988. Detection of induced triploid scallop, Chlamys nobilis by DNA microfluorometry with DAPI staining. Aquaculture 69:201-209.
- 5)Okazaki,T., Tanaka,Y. and Naganuma,H.
1988. A record of the sockeye salmon run to the Tama River, Honshu, Japan. Jpn. J. Ichthyol. 35(1):83-86.
- 6)Fukao,R. and Okazaki,T.
1987. A study on the divergence of Japanese fishes of the genus Neoclinus. Jpn. J. Ichthyol. 34(3):309-323.
- 7)Okazaki,T.
1988. Population structures and homing traits of the fish of the genus Oncorhynchus. Kotaigun seitaigakkaihou 44:31-38.
- 8)Murai,T., Ogata,H., Hirasawa,Y., Akiyama,T. and Nose,T.
1987. Portal absorption and hepatic uptake of amino acids in rainbow trout force-fed complete diets containing casein or crystalline amino acids. Nippon Suisan Gakkaishi 53(10) 1847-1859.
- 9)Hara,A., Ouchi,Y., Nagahama,Y. and Nose,T.
1987. Identification of female-specific serum proteins (vitellogenin) and their related egg yolk protein in red sea bream,Pagrus major. Bull. Natl. Res. Inst. Aquaculture 12:25-35.
- 10)Murai,T., Ogata,H., Kosutarak,P. and Arai,S.
1987. Effects of methanol treatment and amino acid supplementation on utilization of soy flour by chum salmon fingerlings. Bull. Natl. Res. Inst. Aquaculture 12:37-41.
- 11)Suzuki,T.
1988. Ultrastructure of the excretory organs and urinary component of the marine bivalve, Atrina (S.) pectinata. Comp. Biochem. Physiol. 90A(1):79-83.

- 12) Yamaguchi, K. and Mori, K.
1988. Hemagglutinin activities in pearl oyster, Pinctada fucata martensii, hemolymph. Bull. Natl. Res. Inst. Aquaculture 13:25-32.
- 13) Murai, T., Akiyama, T., Ogata, H. and Suzuki, T.
1988. Interaction of dietary oxidized fish oil and glutathion on fingerling yellowtail Seriola quinqueradiata. Nippon Suisan Gakkaishi 54:145-149.
- 14) Shimma, H., Kijima, T. and Maeda, H.
1987. Development of the gonads of one-year-old rainbow trout reared at low in summer and high in water temperature. Bull. Natl. Res. Inst. Aquaculture 12:9-16.
- 15) Shimma, H. and Kitamura, H.
1987. Maturation of female yearling of amago salmon Oncorhynchus rhodurus and acquisition of fertilized eggs (short paper). Bull. Natl. Res. Inst. Aquaculture 12:93-96.
- 16) Maruyama, S. and Konishi, K.
1979. The occurrence of pinnotherid crabs on the northeast coast of Hokkaido, faced on the Okhotsk Sea. Hokusuisi Geppou 36:201-208.
- 17) Hirose, K. and Arai, S.
1988. Effect of LH-RH cholesterol pellet on ovarian maturation in Ayu, Plecoglossus altivelis. Bull. Natl. Res. Inst. Aquaculture 13:11-16.
- 18) Tanaka, H.
1988. Effects of estradiol-17 β on gonadal sex differentiation in flounder, Paralichthys olivaceus. Bull. Natl. Res. Inst. Aquaculture 13:45-53.
- 19) Kagawa, H. and Takano, K.
1979. Ultrastructure and histochemistry of granulosa cells of pre- and post-ovulatory follicles in the ovary of the medaka, Oryzias latipes. Bull. Fac. Fish. Hokkaido. Univ. 30(3):191-204.
- 20) Kagawa, H. and Nagashima, Y.
1980. Ultrastructural localization of $4^5-3\beta$ -hydroxysteroid dehydrogenase in the interrenal cells of the goldfish (Carassius auratus). Cell. Tissue. Res. 212:225-231.
- 21) Nagahama, Y., Kagawa, H. and Tashiro, F.
1980. The In vitro effects of various gonadotropins and steroid hormones on oocyte maturation in amago salmon Oncorhynchus rhodurus and rainbow trout Salmo gairdneri. Bull. Japan. Soc. Sci. Fish. 46(9):1097-1102.
- 22) Kagawa, H. and Nagahama, Y.
1981. In vitro effects of prostaglandins on ovulation in goldfish Carassius auratus. Bull. Japan. Soc. Sci. Fish. 47(9):1119-1121.
- 23) Kagawa, H., Takano, K. and Nagahama, Y.

1981. Correlation of plasma estradiol-17 β and progesterone levels with ultrastructure and histochemistry of ovarian follicles in the white-spotted char, Salvelinus leucomaenis. Cell Tissue Res 218:315-329.
- 24) Nagahama, Y., Kagawa, H. and Young, G.
1982. Cellular sources of sex steroids in teleost gonads. Can. J. Fish. Aquat. Sci. 39:56-64.
- 25) Nagahama, Y. and Kagawa, H.
1982. In vitro steroid production in the postovulatory follicles of the amago salmon, Oncorhynchus rhodurus, in response to salmon gonadotropin. J. Exp. Zool. 219:105-109.
- 26) Kagawa, H., Young, G. and Nagahama, Y.
1982. Estradiol-17 production in isolated amago salmon (Oncorhynchus rhodurus) ovarian follicles and its stimulation by gonadotropins. Gen. Com. Endocrinol 47:361-365.
- 27) Kagawa, H., Young, G. and Nagahama, Y.
1983. Changes in plasma steroid hormone levels during gonadal maturation in female goldfish Carassius auratus. Bull. Japan. Soc. Sci. Fish. 49(12):1783-1787.
- 28) Kagawa, H., Young, G., Adachi, S. and Nagahama, Y.
1982. Estradiol-17 β production in amago salmon (Oncorhynchus rhodurus) ovarian follicles: Role of the thecal and granulosa cells. Gen. Com. Endocrinol. 47:440-448.
- 29) Young, G., Kagawa, H. and Nagahama, Y.
1983. Evidence for a decrease in aromatase activity in the ovarian granulosa cells of amago salmon (Oncorhynchus rhodurus) associated with final oocyte maturation. Biol. Reprod. 29:310-315.
- 30) Kagawa, H., Young, G. and Nagahama, Y.
1983. Relationship between seasonal plasma estradiol-17 β and testosterone levels and In vitro production by ovarian follicles of amago salmon (Oncorhynchus rhodurus). Biol. Reprod. 29:301-309.
- 31) Salmon, C., Kagawa, H., Adachi, S., Nagahama, Y. and Fontaine, Y.
1984. Mise en évidence de sites de liaison spécifique de la gonadotropine de saumon chum (Oncorhynchus rhodurus). C. R. Acad. Sc. Paris, t.298, Serie III, 12:337-340.
- 32) Kagawa, H., Young, G. and Nagahama, Y.
1984. In vitro estradiol-17 β and testosterone production by ovarian follicles of the goldfish, Carassius auratus. Gen. Com. Endocrinol. 54:139-143.
- 33) Kagawa, H.
1985. Ultrastructural and histochemical observations regarding the ovarian follicles of the amago salmon (Oncorhynchus rhodurus). J. Uoeh 7(1):27-35.
- 34) Kagawa, H., Young, G., Adachi, S. and Nagahama, Y.
1985. Estrogen synthesis in the teleost ovarian follicle. The two-cell

- type model in salmonids. In: Salmonid reproduction. (eds. by Iwamoto, R.X And S. Sower) pp 20-25, Washington Sea Grant Program. University of Washington. Washington
- 35) Kanamori, A., Kagawa, H. and Nagahama, Y.
1987. Gonadotropin receptors in the postovulatory ovary of amago salmon (Oncorhynchus rhodurus). Gen. Com. Endocrinol. 66:210-217.
- 36) Nagahama, Y., Young, G., Ueda, H., Kagawa, H. and Adachi, S.
1985. Endocrine control of final gamete maturation in salmonids. In: Salmonid reproduction. (eds. by Iwamoto, R.X And S. Sower) pp ~~20-25~~, 9-19 Washington Sea Grant Program. University of Washington. Washington
- 37) Muraoka, K. and Konishi, K.
1977. Note on the first zoea of Pinnotheres sunensis Shen (Crustacea, brachyura, pinnotheridae) from Tokyo Bay. Research on Crustacea. 8:46-50.
- 38) Konishi, K.
1981. A description of laboratory-reared larvae of the pinnotherid crab Sakaina japonica serene (Decapoda, brachyura). Jour. Fac. Sci. Hokkaido Univ. Ser. VI, 22(3):165-176.
- 39) Iwata, F. and Konishi, K.
1981. larval development in laboratory of Cancer amphioetus Rathbun, in composition with those of seven other species of Cancer (Decapoda, brachyura). Publ. Seto. Mar. Biol. Lab., XXXVI(4/6):369-391.
- 40) Konishi, K.
1981. A description of laboratory-reared larvae of the commensal crab Pinnaxodes mutuensissakai (Decapoda, brachyura) from Hokkaido, Japan. Annotationes Zoologicae Japonenses 54(3):213-229.
- 41) Konishi, K.
1983. Larvae of the Pinnotherid crabs (Crustacea, brachyura) found in the plankton of Osoro Bay, Hokkaido. Jour. Fac. Sci. Hokkaido Univ. Ser. VI, Zool. 23(3):266-295.
- 42) Konishi, K.
1983. A note on classification of brachyuran larvae - Aikawa's zoeal classification. Kaiyou to seibutu 27:256-261.
- 43) Konishi, K.
1985. Laboratory rearing and observation of brachyuran larvae. Seibutsu kyouzai 19:107-116.
- 44) Mawatari, S.F., Konishi, K., Hoshikawa, H., Ishimaru, S., Kuwahara, Y. and Sato, T.
1985. A list of marine invertebrates collected around the Kikonai marine station. 1. Molluscs and amphipod and decapod crustaceans. Seibutsu kyouzai 19:1-10.
- 45) Quintana, R. and Konishi, K.
1986. On the prezoal stage: observations on three Pagurus species (decapoda, anobura). Journal of Natural History 20:837-844.

- 46)Konishi,K.
1986. Larval development of the stone crab, Hapalogaster dentata (De haan,1844) (Crustacea: anomura: lithodidae) reared in the laboratory. Jour. Fac. Sci. Hokkaido Univ. Ser. VI, Zool. 24(3):155-172.
- 47)Konishi,K. and Quintana,R.
1987. Notes on the prezoea stage of decapod crustaceans. Kaiyou to seibutsu 52:372-378.
- 48)Muraoka,K. and Konishi,K.
1987. The first zoeal stage of the porcellanid crab, petrolisthes japonicus(De haan, 1849) with special reference to zoeal features of Petrolisthes(Crustacea: anomura). Research on Crustacea 16:57-65.
- 49)Konishi,K. and Quintana,R.
1987. The larval stage of Pagurus brachiomastus(Thalwitzer, 1892) (Crustacea: anomura) reared in the laboratory. Zool. Sci 4:349-365.
- 50)Konishi,K.
1987. Larval development of the spiny sand crab Lophomastix japonica(Durufle, 1889) (Crustacea, anomura, albuneidae) under laboratory conditions. Publ. Seto. Mar. lab. 32(1/3):123-139.
- 51)Konishi,K.
1987. The larval development of Pachycheles stevensistimpson, 1858 (Crustacea: anomura: porcellanidae) under laboratory conditions. Journal of Crustacean Biology 7(3):481-492.
- 52)Konishi,K. and Quintana,R.
1988. The Larval stages of three pagurid crabs (Crustacea: anomura: paguridae) from Hokkaido, Japan. Zool. Sci. 5:463-482.
- 53)Mawatari,S.F., Kubota,S., Konishi,K., Sato,T., Kuwahara,Y., Yamada,M.
1988. A list of marine invertebrates collected around the Kikonai marine station. 2. Hydrozoa, Bryozoa, Pycnogonida and Decapoda, and Gastropoda and Bivalvia. Seibutu kyoza 23:1-12.
- 54)Sakaguchi,S., Hara,T., Matsusato,T. Shibahara,T., Yamagata,Y., Kawai,H. and Maeno,Y.
1987. Scoliosis of cultured yellowtail caused by parasitic Myxobolus buri. Bull. Natl. Res. Inst. Aquaculture 12:79-86.
- 55)Sakaguchi,S., Yamagata,Y. and Sako,H.
1987. Reidentification of Philometra parasitic on the red sea bream. Bull. Natl. Res. Inst. Aquaculture 12:69-72.
- 56)Sakaguchi,S., Shibahara,T. and Yamagata,Y.
1987. Parasitic ecology of a Philometra lateolabracis parasite on the red sea bream. Bull. Natl. Res. Inst. Aquaculture 12:73-78.
- 57)Sorimachi,M. and Egusa,S.
1987. A histopathological study of ICDV infection of Japanese eel, Anguilla japonica. Bull. Natl. Res. Inst. Aquaculture 12:87-92.

- 58)Sako,H.
1988. Survival of fish pathogenic bacteria in seawater. Bull. Natl. Res. Inst. Aquaculture 13:45-53.
- 59)Miwa,S., Tagawa,M., Inui,Y. and Hirano,T.
1988. Thyroxine surge in metamorphosing flounder larvae. Gen. Comp. Endocrinol. 70:158-163.
- 60)Miwa,S. and Inui,Y.
1987. Effects of various doses of thyroxine and triiodothyronine on the metamorphosis of flounder(Paralichthys olivaceus). Bull. Natl. Res. Inst. Aquaculture 12:356-363.
- 61)Nakanishi,T. and Onozato,H.
1987. Variability in the growth of isogenic crucian carp, Carassius gibelio langsdorfii. Nippon Suisan Gakkaishi 53(12):2099-2104.
- 62)Nakanishi,T.
1987. Transferability of immune plasma and pronephric cells in isogenic, allogeneic and xenogeneic transfer system in crucian carp. Develop. Comp. Immunol. 11:521-528.
- 63)Nakanishi,T.
1987. Histocompatibility analyses in tetraploids induced from clonal triploid crucian carp and in gynogenetic diploid goldfish. J. Fish. Biol. 31(Suppl.A):35-40.
- 64)Ikeda,K.
1987. Effects of heat treatment on agglutinin activity in serum of kokanee(Oncorhynchus nerka). Bull. Natl. Res. Inst. Aquaculture 12:61-64.
- 65)Ikeda,K.
1987. Heat resistances of partially purified antibodies of kokanee, amago salmon and rabbit. Bull. Natl. Res. Inst. Aquaculture 12:65-68.
- 66)Ikeda,K., Kumada,H. and Shmima,Y.
1987. Protein binding states of zinc, cadmium and magnesium in serum of coho salmon(Oncorhynchus kisutch). Bull. Natl. Res. Inst. Aquaculture 12:43-52.
- 67)Imai,I., Itoh,K. and Anraku,M.
1984. Extinction dilution method for enumeration of dormant cells of red tide organisms in Marine sediments. Bull. Plankton Soc. Japan 31(2):123-124.
- 68)Honjo,T.
1985. Glycocalyx on the cell surface in raphidophycean algae. Suisan no kenkyu 18:14-15.
- 69)Honjo,T.
1986. Growth potential in wild phytoplankton.
In: A manual for environmental research of marine coast(Ed. by Japan Oceanographic Society), pp.168-172. Koseisha koseikaku, Tokyo.

- 70) Honjo, T.
1987. Growth Potential of Chattonella marina (Raphidophyceae) collected in Gokasho Bay, central Japan. *Bull. Plankt. Soc. Japan*, 34(2):119-124.
- 71) Honjo, T.
1987. On raphidophycean Heterosigma akashiwo and dinophycean Gymnodinium nagasakiense. *Science of Red Tide* (ed. by T. Okaichi), pp.219-237. Koseisysha, Tokyo.
- 72) Matsushima, T., Kitamura, S., Takei, K., Satou, M., Okumoto, N. and Ueda, K.
1986. An electromyographic study of sexual behavior in Hime Salmon (Landlocked sockeye salmon, Oncorhynchus nerka). *Zool. Sci.* 3:563-567.
- 73) Kitamura, S.
1987. Electroretinogram (ERG) in the rock fish, Sebastes marmoratus-I Relationship between stimulus and responses in the dark adapted state. *Bull. Natl. Res. Inst. Aquaculture* 12:53-60.
- 74) Fujii, K., Hara, A., Hirose, K. and Maruyama, T.
1987. Immunological and electrophoretical studies of specific serum protein induced by estrogen treatment in hybrid sturgeon between female Huso huso and male Acipenser ruthenus, so-called "Bester". *Bull. Natl. Res. Inst. Aquaculture* 12:17-24.
- 75) Nakanishi, T.
1987. Chromosomal manipulation in fish. *Sigen Technology* 226:13-17.
- 76) Machii, A., Wada, K.T., Awaji, M., Nakamura, H.K. And Townsley S.J.
1988. Some characters of cells of the midgut and chytrids from primary culture of the prawn Penaeus japonicus. *Invertebrate and Fish Tissue Culture* (eds. by Y. Kuroda, E. Kurstak and K. Maramorosch), pp. 11-14. Japan Scientific Society Press Tokyo/Springer-Verlag, Berlin.
- 77) Konishi, K.
1977. On the three species of the pinnotherid crabs from Hokkaido, Japan. *Jour. Fac. Sci. Hokkaido Univ. Ser. VI, Zool.* 20:605-611.
- 78) Tanaka, N.
1987. Attaching diatoms as food for bivalves and their cultivation. *Suisan doboku* 24(1):37-41.
- 79) Tanaka, N.
1987. mass-culture of attaching diatoms and their usefulness. *Monthly Report of Japan Fish. Resources Conservation Ass.* 279:9-16.
- 80) Tanaka, N. and Asakawa, A.
1988. Allelopathy in seaweeds. *Kagaku to seibutsu* 26(2):71-73.
- 81) Tanaka, N. and Ohwada, K.
1988. Decomposition of agar, protein, and organic phosphate by marine epiphytic diatoms. *Nippon Suisan Gakkaishi*. 54(4):725-727.

- 82)Asakawa,A., Ohwada,K. and Tanaka,N.
1988. The seasonal variation in chemical composition of Eiscnia bicyclis and Ecklonia cava, collected from the coast of shima peninsula. Bull. Natl. Res. Inst. Aquaculture 13:33-44.
- 83)Ihkura,T.
1987. Effect of artificial substrate on the cultivation of phytal animals. Fisheries Engineering, (24)1.
- 84)Toda,S. and Ihkura,T.
1988. Observation of micro environment in Sargassum forest. Progress report of Marine plant for 1987. pp.1-16.
- 85)Kakuno,A. and Kimura,S.
1987. Acute toxicity of Bis (tributyltin) oxide to girella (Girella punctata). Bull. Tokai. Reg. Lab. 123:41-44.
- 86)Umebayashi,O.
1988. Individual leaf growth of vegetative shoot of elgrass Zostera marina. Bull. Tokai. Reg. Lab. 125:1-13.
- 87)Shimizu,A. and Kimura,S.
1987. Effect of Bis (tributyltin) oxide on gonadal development of a salt-water goby, Chasmichthys dolicognathus : Expose durling maturing period. Bull. Tokai. Reg. Fish. Res. Lab. 123:45-49.
- 88)Yamada,H.
1987. Water pollution by organophosphoric acid triesters and its effects on aquatic organisms-A review. Bull. Tokai. Reg. Fish. Res. Lab. 123:15-30.
- 89)Akiyama,K., Kodaka,G. and Hara,M.
1988. Genetic differences between natural and cultured populations of Porphyra yezoensis. Tohoku Journal of Agricultural Research 38(1-4):27-34.
- 90)Fujio,Y., Tanaka,M.Y. and Hara,M.
1987. Enzyme polymorphism and population structure of the haploid laver Porphyra yezoensis. Nippon Suisan Gakkaishi 53(3):357-362.
- 91)Fujio,Y., Kodaka,P.L.G. and Hara,M.
1988. Electrophoretic variants characterstic heterozygetes in haploid laver Porphyra sp. Nippon Suisan Gakkaishi 54(6):969-974.
- 92)Hara,M. and Ishikawa,Y.
1988. Difference of the growth and shapes among four local forms, Undaria pinnatifida, cultured in same grounds. Fish Genetics and Breeding Science 13:29-33.
- 93)Kawamura,T., Nimura,Y. and Hirano,R.
1988. Effects of bacterial films on diatom attachment in the initial phase of marine fouling. J. Oceanographical Soc. of Japan 44:1-5.

- 94)Kurata,K., S Suzuki,M., Shiraishi,K. and Taniguchi,K.
1988. Spatane-type diterpenes with biological activity from the brown alga Dilophus okanurai. *Phytochemistry* 27(5):1321-1324.
- 95)Taniguchi,K.
1988. Marine afforestation of eisenia bicyetis (Laminariaceae; Pheophyta). *Kaiyoukagaku* 20(6)363-368.
- 96)Taniguchi,K. and Yamada,H.
1988. Annual variation and productivity of the Sargassum horneri population in Matsushima Bay on the pacific coast of Japan. *Bull. Tohoku. Reg. Fish. Res. Lab* 50:59-65.
- 97)Fukuhara,O. and Fushimi,T.
1987. Fin differentiation and squamation of artificially reared grouper, Epinephelus akaara. *Aquaculture*, 69:379-386.
- 98)Imai,I. and Itho,K.
1984. Distribution of heterotrophics microflagellates in Suo-Nada, western seto inland sea, in May 1983. *Bull. Nansei. Reg. Fish. Res. Lab.* 17:219-231.
- 99)Imai,I.
1984. Size distribution and biomass of bacteria in Suo-Nada, western seto inland sea. *Bull. Nansei. Reg. Fish. Res. Lab.* 17:183-196.
- 100)Imai,I., Itoh,K. and Anraku,M.
1984. Distribution of dormant cells of Chattonella in Harima-Nada, eastern seto inland sea, and temperature characteristics of germination. *Bull. Plankton. Soc. Japan.* 31(1):35-42.
- 101)Imai,I. and Itoh,K.
1985. Distribution of dormant cells of Chattonella in bottom sediments of Harima-Nada, eastern seto inland sea, in April, 1984. *Bull. Nansei. Reg. Fish. Res. Lab.* 19:43-52.
- 102)Imai,I. and Itoh,K.
1986. A preliminary note on the cysts of Chattonella(Raphidophyceae), red tide flagellates, found in bottom sediment in Suo-Nada, western seto inland sea, Japan. *Bull. Plakton. Soc. Japan.* 33(1):61-63.
- 103)Imai,I., Itoh,K., Terada,K. and Kamizono,M.
1986. Distribution of dormant cells of Chattonella(Raphidophyceae) and occurrence of summer red tide in Suo-Nada, western seto inland sea. *Bull. Japan. Soc. Sci. Fish.* 52(9):1665-1671.
- 104)Imai,I.
1987. Size distribution, number and biomass of bacteria in intertidal sediments and seawater of Ohmi Bay, Japan. *Bull. Jap. Soc. Microb. Ecol. Nihon Biseibutsu Seitai Gakkaiho* 2(1):1-11.
- 105)Imai,I. and Itoh,K.
1987. Annual life cycle of Chattonella spp., causative flagellates of noxious red tides in the inland sea of Japan. *Marine Biology.* 94:287-292.

- 106) Ishioka, H., Hatakeyama, Y. and Sakaguchi, S.
1988. The hearing ability of the red sea bream Pagrus major. Nippon Suisan Gakkaishi 54(6):947-951.
- 107) Itoh, K. and Imai, I.
1986. Grazing on Chattonella marina and production of fecal pellets by copepods. Bull. Nansei. Reg. Fish. Res. Lab. 20:115-126.
- 108) Itoh, K. and Imai, I.
1988. Recent advances in preventive strategies and the occurrence of red tide. Studies on Fisheries Oceanography. pp 228-237. (ed by the Japanese Society of Fisheries Oceanography.)
- 109) Kamiyama, T. and Aizawa, Y.
1987. Growth characteristics of two tintinnid ciliates, tintinnopsis beroidea and Amphorella quadrilineata, in laboratory cultures. Bull. Plankton. Soc. Japan. 34(2):185-191.
- 110) Kishida, T.
1988. Vertical and horizontal distribution of eggs and larvae of Japanese spanish mackerel in the central water of the Seto inland sea. Nippon Suisan Gakkaishi. 54(1)1-8.
- 111) Masaki, Y., Ito, H., Kamijyo, Y., Yokomatsu, Y., Ogawa, H., Yamaguchi, Y. and Tokai, T.
1987. Sexual maturity and spawning season of Japanese flounder in Suo-Nada. Nippon Suisan Gakkaishi 53(7):1181-1190.
- 112) Matsuo, Y.
1986. Vertical distribution of chlorophyll-a and phaeopigments on an intertidal flat in Ohmi Bay. Bull. Nansei. Reg. Fish. Res. Lab. 20:127-136.
- 113) Ouchi K., Adachi, S. and Nagahama, Y.
1988. Changes in plasma levels of steroid hormones during sexual maturation of female red seabream Pagrus major. Nippon Suisan Gakkaishi 54(4):585-591.
- 114) Tanaka, K.
1988. Study on the abalone culture in the coast of Awa region, Chiba prefecture. Bull. Jap. Sea. Reg. Fish. Res. Lab. 38:21-132.
- 115) Yamaguchi, M. and Anraku, M.
1984. Primary productivity in Suo-Nada, western Seto inland sea. Bull. Nansei. Reg. Fish. Res. Lab. 17:135-149.
- 116) Matsuoka, M.
1987. Development of the skeletal tissues and skeletal muscles in the red sea bream. Bull. Seikai Reg. Fish. Res. Lab. 65:1-114.
- 117) Morioka, Y., Nakashima, J. and Kimoto, K.
1987. Oxygen consumption and biological productivity of mysid population in a small inlet of Kyushu. Bull. Seikai Reg. Fish. Res. Lab. 65:115-123.