

1991 LITERATURE EXCHANGE FROM THE JAPANESE AQUACULTURE PANEL, UJNR,
TO THE UNITED STATES PANEL, UJNR

Prepared by:
Masahiko Awaji
National Research Institute
of Aquaculture
Fisheries Agency

Hokkaido National Fisheries Research Institute

1. Goto, A. and T. Andoh (1990) Genetic divergence between the sibling species of river-sculpin, *Cottus amblystomopsis* and *C. nozawae*, with special reference to speciation. *Env. Biol. Fish.*, 28, 257-266.
2. Nabata, S. (1991) List of marine algae from the Shiribeshi coast of Hokkaido. *Sci. Rep. Hokkaido Fish. Exp. Stn.*, 36, 19-38.
3. Saitoh, S., T. Mori, M. Kusakari and K. Miura (1991) Studies on the spawning of Japanese flounder (*Paralichthys olivaceus*). I. Induction of spawning of Japanese flounder by manipulation of water temperature. *Sci. Rep. Hokkaido Fish. Exp. Stn.*, 36, 71-79.
4. Sakanishi, Y. and N. Saga (1991) Evaluation of the UV sensitivity of the brown alga *Macrocystis pyrifera* by a vital-staining method. *Bull. Hokkaido Natl. Fish. Res. Inst.*, 55, 209-212.
5. Sakata, K., Y. Iwase, K. Kato, K. Ina and Y. Machiguchi (1991) A simple feeding inhibitor assay for marine herbivorous gastropods and the sea urchin *Strongylocentrotus intermedius* and its application to unpalatable algal extracts. *Nippon Suisan Gakkaishi*, 57(2), 261-265.
6. Sakata, K., K. Kato, Y. Iwase, H. Okada, K. Ina and Y. Machiguchi (1991) Feeding-stimulant activity of algal glycerolipids for marine herbivorous gastropods. *J. Chem. Ecol.*, 17, 185-193.
7. Sakurai, I., T. Miyamoto and K. Takahashi (1991) Environmental characteristics and distribution of young bivalves in Japanese surf clam bed off Tomakomai coast, Hokkaido. *Sci. Rep. Hokkaido Fish. Exp. Stn.*, 36, 39-59.
8. Tajima, K., K. Yamashita and M. Fukuchi (1991) Studies on the artificial seed production of the sea urchin, *Strongylocentrotus intermedius*. II. The collection of metamorphosed larvae. *Sci. Rep. Hokkaido Fish. Exp. Stn.*, 36, 61-70.

Hokkaido Salmon Hatchery

9. Ban, M. and K. Yamauchi (1991) Seasonal changes in seawater

- adaptability of the hatchery reared juvenile sockeye salmon, *Oncorhynchus nerka*. Sci. Rep. Hokkaido Salmon Hatchery, 45, 25-33.
10. Ishida, T., S. Hashimoto and Y. Takada (1991) Planktonic and benthic crustaceans of lake Kussharo, in 1990. Sci. Rep. Hokkaido Salmon Hatchery, 45, 61-64.
 11. Kaeriyama, M. (1991) Dynamics of the lacustrine sockeye salmon population in lake Shikotsu, Hokkaido. Sci. Rep. Hokkaido Salmon Hatchery, 45, 1-24.
 12. Kaeriyama, M. (1991) Development and growth of the genus *Oncorhynchus*. 3. Growth curve. Tech. Rep. Hokkaido Salmon Hatchery, 160, 47-52.
 13. Kusakari, M. and S. Urawa (1990) Histopathology of the skin of yearling Japanese flounder *Paralichthys olivaceus* infected with the flagellate *Ichthyobodo* sp. Fish Pathology, 25(2), 59-68.
 14. Mayama, H. (1990) Masu salmon propagation in Hokkaido, Japan. Bull. Inst. Zool., Academia Sinica, 29(3, Supplement), 95-104.
 15. Mayama, H. (1991) Notes on the life of salmonid fishes in the Nishibetsu River, eastern Hokkaido. Tech. Rep. Hokkaido Salmon Hatchery, 160, 11-23.
 16. Nagasawa, K. and S. Urawa (1991) New records of the parasitic copepod *Salmincola stellatus* from Sakhalin taimen (*Hucho perryi*) in Hokkaido, with a note on its attachment site. Sci. Rep. Hokkaido Salmon Hatchery, 45, 57-59.
 17. Ohkuma, K. (1991) Differences in scale characteristics taken from various parts of masu salmon, *Oncorhynchus masou* (Brevoort) and its scale development. Sci. Rep. Hokkaido Salmon Hatchery, 45, 35-46.
 18. Seki, J. and I. Shimizu (1991) Characteristics of an oligotrophic lake in Hokkaido - II. Annual and seasonal variations in the biomass of zooplankton. in "Limnological comparison of characteristics of water quality in Chinese and Japanese lakes", eds. M. Aizaki and H. Tachibana, Nissei Eblo, Tsukuba, 149-151.
 19. Shimizu, I. and J. Seki (1991) Characteristics of an oligotrophic lake in Hokkaido - I. Chemical characteristics of lake Shikotsu. in "Limnological comparison of characteristics of

- water quality in Chinese and Japanese lakes", eds. M.Aizaki and H.Tachibana, Nissei Eblo, Tsukuba, 141-147.
20. Shimizu, I and M.Kaeriyama (1991) Biological and chemical characteristics of chum salmon during their spawning migration period. Sci. Rep. Hokkaido Salmon Hatchery, 45, 47-56.
 21. Ueda, H., O.Hiroi, K.Yamauchi, A.Hara, H.Kagawa, S.Adachi and Y.Nagahama (1991) Changes in serum steroid hormone levels and *in vitro* steroid hormone production in the ovaries of female chum salmon *Oncorhynchus keta* during spawning migration. Nippon Suisan Gakkaishi, 57(5), 881-884.
 22. Ueda, H., O.Hiroi, K.Yamauchi, A.Hara, H.Kagawa, S.Adachi and Y.Nagahama (1991) Changes in serum steroid hormone levels and *in vitro* steroid hormone production in the testes of male chum salmon *Oncorhynchus keta* during spawning migration. Nippon Suisan Gakkaishi, 57(6), 1111-1114.
 23. Urawa, S. and J.R.Arthur (1991) First record of the parasitic ciliate *Trichodina truttae* Mueller, 1937 on chum salmon fry (*Oncorhynchus keta*) from Japan. Gyobyo Kenkyu, 26(2), 83-89.
 24. Urawa, S., K.Muroga and S.Kasahara (1991) Growth and fecundity of the parasitic copepod *Neoergasilus japonicus* (Ergasilidae). Proc. Fourth Int. Conf. Copepoda; Bull. Plankton Soc. Japan, Spec. Vol., 619-625.
 25. Urawa, S. (1991) A review of sockeye salmon production in the Nishibetsu river in eastern Hokkaido, Japan. Tech. Rep. Hokkaido Salmon Hatchery, 160, 3-10.
 26. Bibliography of salmonids published in Japan (4) 1989. (1991) eds. S.Urawa and T.Azuma, Tech. Rep. Hokkaido Salmon Hatchery, 160, 67-90.
 27. Abstracts of papers presented at the 15th salmon seminar of the Hokkaido Salmon Hatchery. (1991) Tech. Rep. Hokkaido Salmon Hatchery, 160, 59-66.

Tohoku National Fisheries Research Institute

28. Kawaguchi, K., Y.Yamashita and A.Hayashi (1990) Some aspects of spawning of the reared Japanese anchovy (*Engraulis japonicus* H.) in relation to the photoperiod, water temperature and

- starvation. Bull. Japan. Soc. Fish. Oceanogr., 54(4), 364-372.
29. Kishi, M.J., S. Kimura, H. Nakata and Y. Yamashita (1991) A biomass-based model for the sand lance (*Ammodytes personatus*) in Seto inland sea, Japan. Ecological Modelling, 54, 247-263.
30. Kurata, K., K. Taniguchi, K. Shiraishi and M. Suzuki (1990) Feeding-deterrent diterpenes from the brown alga *Dilophus okamurai*. Phytochemistry, 29(11), 3453-3455.
31. Kurata, K., K. Taniguchi, K. Shiraishi and M. Suzuki (1990) A C₂₆ sterol from the brown alga *Eisenia bicyclis*. Phytochemistry, 29(11), 3678-3680.
32. Taniguchi, K. (1991) Marine algal recolonization on the denuded sublittoral rock surfaces off Oshika peninsula, Japan. Bull. Tohoku Natl. Fish. Res. Inst., 53, 1-5.
33. Taniguchi, K., K. Isogami and H. Kojima (1991) Observations on the growth and maturation of 2-4 year old plants of *Eisenia bicyclis* (Laminariaceae, Phaeophyta). Jpn. J. Phycol., 39, 43-47.

National Research Institute of Fisheries Engineering

34. Hagino, S. (1991) Fishing effectiveness of the artificial reef in Japan. in "Proceedings of the Japan-U.S. Symposium on Artificial Habitats for Fisheries", Japan International Marine Science and Technology Federation, 63-67.
35. Higano, J. (1991) Artificial habitats for bivalves of sandy beach and estuary. in "Proceedings of the Japan-U.S. Symposium on Artificial Habitats for Fisheries", Japan International Marine Science and Technology Federation, 106-111.
36. Kimoto, K. (1991) Swarming behavior of copepod, and its significance for predator-prey relationships between juvenile red sea bream and copepods. Rept. Fish. Res. Invest. Japan. Gov., 27, 51-60.
37. Yasunaga, Y. and J. Higano (1991) Ethological experiments of relation between gathering of fish to the fish reef area and physico-chemical environmental factors. Tech. Rep. Natl. Res. Inst. Fish. Engineering, 13, 1-13.

National Research Institute of Fisheries Science

38. Iguchi, K. and N. Mizuno (1990) Diel changes of larval drift among amphidromous gobies in Japan, especially *Rhinogobius brunneus*. J. Fish Biol., 37, 255-264.
39. Iguchi, K. (1991) Visual stimuli causing the aggressive behaviour of Ayu *Plecoglossus altivelis*. Bull. Natl. Res. Inst. Fish. Sci., 2, 15-23.
40. Iguchi, K. and N. Mizuno (1991) Mechanisms of embryonic drift in the amphidromous goby, *Rhinogobius brunneus*. Env. Biol. Fish., 31, 295-300.
41. Ishida, N. (1991) Concentrations of oxolinic acid and its metabolites in the urine of rainbow trout. Nippon Suisan Gakkaishi, 57(1), 179.
42. Kiso, K. (1990) Polymorphism of life form in masu salmon (*Oncorhynchus masou*) in the rivers of southern Sanriku district, Honshu, Japan. Bull. Inst. Zool., Academia Sinica, 29(3, Suppl.), 27-39.
43. Yokoyama, M. and J. Nakazoe (1991) Effects of dietary protein levels on free amino acid and glutathione contents in the tissues of rainbow trout. Comp. Biochem. Physiol., 99A(1/2), 203-206.

National Research Institute of Far Seas Fisheries

44. Nagasawa, K., J. Bresciani and J. Lützen (1991) *Ischnochitonika japonica*, new species (Copepoda), a parasite on *Ischnochiton* (*Ischnoradsia*) *hakodadensis* (Pilsbry) (Polyplacophora: Ischnochitonidae) from the sea of Japan. J. Crustacean Biol., 11(2), 315-321.
45. Nagasawa, K., K. Takahashi, S. Tanaka and M. Nagata (1991) Ecology of *Pectenophilus ornatus*, a copepod parasite of the Japanese scallop *Patinopecten yessoensis*. Proceedings of the Fourth International Conference on Copepoda; Bull. Plankton Soc. Japan, Spec. Vol., 495-502.
46. Ogura, M., Y. Ishida and S. Ito (1991) Growth variation of coho salmon *Oncorhynchus kisutch* in the western North Pacific.

Nippon Suisan Gakkaishi, 57(6), 1089-1093.

47. Ueno, Y. and K. Hiramatsu (1990) A method for estimating mortality coefficients of chum salmon *Oncorhynchus keta* from tagging experiments. Nippon Suisan Gakkaishi, 56(7), 1079-1082.

National Research Institute of Aquaculture

48. Suzuki, T. (1990) Electrophoretic and chromatographic patterns of hemolymph proteins of the pearl oyster, *Pinctada fucata*. Bull. Natl. Res. Inst. Aquaculture, 18, 7-12.
49. Suzuki, T. and K. Mori (1991) Immunolocalization and *in vitro* secretion of hemolymph lectin of the pearl oyster, *Pinctada fucata martensii*. Zool. Sci., 8(1), 23-29.
50. Yoshinaka, R., S. Mizuta, T. Suzuki and M. Sato (1991) The immunological study on the effect of pepsin digestion on genetically distinct types of collagen in muscle of kuruma prawn *Penaeus japonicus*. Comp. Biochem. Physiol., 98B(1), 59-65.
51. Komaru, A., H. Matsuda, T. Yamakawa and K. T. Wada (1990) Chromosome behavior of meiosis-inhibited eggs with cytochalasin B in Japanese pearl oyster. Nippon Suisan Gakkaishi, 56(9), 1419-1422.
52. Durand, P., K. T. Wada and A. Komaru (1990) Triploidy induction by caffeine-heat shock treatments in the Japanese pearl oyster *Pinctada fucata martensii*. Nippon Suisan Gakkaishi, 56(9), 1423-1425.
53. Wada, K. T. and A. Komaru (1990) Inheritance of white coloration of the prismatic layer of shells in the Japanese pearl oyster *Pinctada fucata martensii* and its importance in the pearl culture industry. Nippon Suisan Gakkaishi, 56(11), 1787-1790.
54. Wada, K. T., J. Scarpa and S. K. Allen, Jr. (1990) Karyotype of the dwarf surfclam *Mulinia lateralis* (Say, 1822) (Mactridae, Bivalvia). J. Shellfish Res., 9, 279-281.
55. Wada, K. T. (1991) The pearl oyster, *Pinctada fucata* (Gould) (Family Pteriidae). in "Estuarine and Marine Bivalve Mollusk Culture" ed. W. Menzel, CRC Press, Boca Raton, Florida, 245-260.

56. Nagoya, H., T. Kimoto and H. Onozato (1990) Diploid gynogenesis induced by suppression of the second or the third cleavage in the goldfish, *Carassius auratus*. Bull. Natl. Res. Inst. Aquaculture, 18, 1-6.
57. Konishi, Y., H. Onozato, A. Yamamoto and K. Fujimori (1991) Characteristics of lysosomal hydrolytic pathway of glycogen in striated muscle of diploid and triploid masu salmon. Nippon Suisan Gakkaishi, 57(5), 943-949.
58. Konishi, Y., H. Onozato, A. Yamamoto and K. Fujimori (1991) Glycogenosomes in hepatocyte of diploid and triploid masu salmon (*Oncorhynchus masou*), mimicking cytology of glycogen storage disease type II. Agric. Biol. Chem., 55(4), 1147-1149.
59. Okazaki, T. (1990) Population structure of masu salmon, *Oncorhynchus masou*, in the species of the genus *Oncorhynchus*. Bull. Inst. Zool., Academia Sinica, 29 (3, Suppl.), 17-25.
60. Okazaki, T. (1991) Genetic population structure in the species of the genus *Oncorhynchus* and fluctuation of its abundance. Rept. Fish. Res. Invest. Japan. Gov., 27, 17-30.
61. Oohara, I., T. Akiyama, H. Aono and K. Mori (1991) Distribution and inter-organic correlations of lysozyme activity in the juvenile bluefin tuna, *Thunnus thynnus*. Bull. Natl. Res. Inst. Aquaculture, 19, 17-26.
62. Hirose, K., H. Kagawa, M. Yoshida, M. Kumakura and H. Yamanaka (1990) Application of LHRH copolymer pellet for induction of final oocyte maturation and ovulation in Ayu *Plecoglossus altivelis*. Nippon Suisan Gakkaishi, 56(11), 1731-1734.
63. Adachi, S., H. Kajiura, H. Kagawa and Y. Nagahama (1990) Identification of aromatizable androgens secreted by isolated thecal cell layers from vitellogenic ovarian follicles of a teleost, amago salmon (*Oncorhynchus rhodurus*). Biomedical Res., 11(5), 359-363.
64. Kagawa, H. (1991) Ultrastructural observations on the ovarian follicles of the yellowtail (*Seriola quinqueradiata*) during oocyte maturation and ovulation. Bull. Natl. Res. Inst. Aquaculture, 19, 1-10.
65. Kagawa, H., H. Tanaka, K. Okuzawa, M. Matsuyama and K. Hirose (1991) Diurnal changes in plasma 17α , 20β -dihydroxy-4-pregnen-3-one levels during spawning season in the red sea bream *Pagrus*

- major*. Nippon Suisan Gakkaishi, 57(4), 769.
66. Konishi, K. and H. Sekiguchi (1990) First-stage phyllosoma of *Scyllarus cultrifer* (Ortmann) (Decapoda, Scyllaridae). Bull. Plankton Soc. Japan, 37(1), 77-82.
 67. Konishi, K. and M. Minagawa (1990) The first zoeal larva of the gulfweed crab *Planes cyaneus* Dana, 1851 (Crustacea: Brachyura: Grapsidae). Proc. Japan. Soc. Syst. Zool., 42, 14-20.
 68. Okuzawa, K., M. Amano, M. Kobayashi, K. Aida, I. Hanyu, Y. Hasegawa and K. Miyamoto (1990) Differences in salmon GnRH and chicken GnRH-II contents in discrete brain areas of male and female rainbow trout according to age and stage of maturity. Gen. Comp. Endocrinol., 80, 116-126.
 69. Okuzawa, K., K. Furukawa, K. Aida and I. Hanyu (1991) Photoperiodic responses of the cyprinid honmoroko *Gnathopogon caerulescens* during the nonbreeding season. Bull. Natl. Res. Inst. Aquaculture, 19, 11-16.
 70. Nakajima, K., C. H. Kweon, S. H. An, T. Shibayama, N. Wakamiya, S. Kato and K. Hirai (1990) Identification of the cross-reactive antigens between Marek's disease virus and pseudorabies virus by monoclonal antibodies. Avian Diseases, 34, 479-484.
 71. Nakajima, K., T. Shibayama, S. Kato, T. Kurimura and K. Hirai (1990) Synthesis and processing of a gp28/32 membrane glycoprotein induced by Marek's disease virus serotype 2. J. Gen. Virology, 71, 1807-1810.
 72. Naito, M., K. Nakajima, S. Osafune, K. Ono and N. Iwa (1991) Detection of Marek's disease virus-specific phosphorylated protein antigen in the feather follicle epitheliums of chickens injected with HVT vaccine. Research Reports of Himeji college of Hyogo, 36, 223-228.
 73. Wang, D. Q., Y. Fukui, T. Ito, K. Nakajima, S. Kato, M. Naiki, T. Kurimura and N. Wakamiya (1990) Heterogeneity of Hanganutziu-Deicher antigen glycoproteins in different species animal sera. Jpn. J. Vet. Sci., 52(3), 567-572.
 74. Ito, T., D. Q. Wang, M. Maru, K. Nakajima, S. Kato, T. Kurimura and N. Wakamiya (1990) Antitumor efficacy of vaccinia virus -modified tumor cell vaccine. Cancer Research, 50, 6915-6918.
 75. Matsumoto, T., O. Yamada, S. Kageyama, N. Wakamiya, K. Nakajima, K. Imai, D. Q. Wang, Y. S. Shin, T. Kurimura, M. Honda, T. Tokunaga,

- M.Ito, T.Meguro and K.Yamada (1990) A longitudinal study of 355 HIV seropositive individuals to evaluate the HIV isolation as an indicator of prognosis. AIDS Research Newsletter, Proceedings of the 4th Congress of the Japanese Society for AIDS Research, 18-19 July 1990, Yokohama, Japanese Society for AIDS Research.
76. Nakanishi, T. (1991) Ontogeny of the immune system in *Sebastiscus marmoratus*: histogenesis of the lymphoid organs and effects of thymectomy. Environmental Biol. Fish., 30, 135-145.
77. Nakanishi, T. and H.Onozato (1990) Clonal ginbuna crucian carp as a model for the study of fish immunology and genetics. NOAA Technical Report NMFS, 92, 45-51.
78. Fuda, H., T.Nakanishi and F.Yamazaki (1991) Preliminary observations on ontogenetic development of the alloimmune reactivity in chum salmon, *Oncorhynchus keta*. Bull. Fac. Fish., Hokkaido Univ., 42(2), 35-38.
79. Hashimoto, K., T.Nakanishi and Y.Kurosawa (1990) Isolation of carp genes encoding major histocompatibility complex antigens. Proc. Natl. Acad. Sci. USA, 87, 6863-6867.
80. Sorimachi, M. (1990) Viral diseases of marine fishes in Japan. Proc. ROC-Japan Symp. Fish Dis., 120-127.
81. Iida, Y., T.Nakai, M.Sorimachi and K.Masumura (1991) Histopathology of a herpesvirus infection in larvae of Japanese flounder *Paralichthys olivaceus*. Dis. aquat. Org., 10, 59-63.
82. Meguro, Y., T.Nakai, K.Muroga and M.Sorimachi (1991) A cell line derived from the fin of Japanese flounder, *Paralichthys olivaceus*. Gyobyo Kenkyu, 26(2), 69-75.
83. Yamano, K., S.Miwa, T.Obinata and Y.Inui (1991) Thyroid hormone regulates developmental changes in muscle during flounder metamorphosis. Gen. Comp. Endocrinol., 81, 464-472.
84. Fujii, K., K.Hirose, A.Hara, M.Shiraishi and T.Maruyama (1991) Use of vitellogenin level as a maturational indicator for artificial spawning of cultured hybrid sturgeon, *Huso huso* X *Acipenser ruthenus*. in "ACIPENSER sturgeon, Proceedings of the 1st international Bordeaux Symposium 1989", ed. P.Williot, Cemagref publ., France, 381-388.
85. Maeda, M. (1991) Microorganisms as bio-piscimedcines for the

growth of fishes and protection from the diseases. in "Microbiological Agents in Agriculture and Fisheries", eds. S.Egusa, H.Komada and H.Iwahana, CMC Publication, Tokyo, 178-193.

86. Maeda, M. and A.Hino (1991) Environmental management for mass culture of the rotifer, *Brachionus plicatilis*. in " Rotifer and Microalgae Culture Systems", eds. W.Fulks and K.L.Main, Oceanic Institute, Honolulu, 125-133.
87. Sugiyama, M., H.Tanaka and K.Fukusho (1991) Toxicity of total ammonia and nitrite nitrogen to young yellowtail, *Seriola quinqueradiata*. Bull. Natl. Res. Inst. Aquaculture, 19, 31-33.
88. Hirano, T., T.Ogasawara, S.Hasegawa, M.Iwata and Y.Nagahama (1990) Changes in plasma hormone levels during loss of hypoosmoregulatory capacity in mature chum salmon (*Oncorhynchus keta*) kept in seawater. Gen. Comp. Endocrinol., 78, 254-262.
89. Iwata, M., T.Shimoyama, N.Sakai, K.Suzuki, H.Ida, K.Muto and U.Akutsu (1990) Effects of sulfuric acid stress on osmoregulatory ability of Japanese char (*Salvelinus leucomaenis*). Bull. Natl. Res. Inst. Aquaculture, 18, 31-37.
90. Sakamoto, T., M.Iwata and T.Hirano (1991) Kinetic studies of growth hormone and prolactin during adaptation of coho salmon, *Oncorhynchus kisutch*, to different salinities. Gen. Comp. Endocrinol., 82, 184-191.
91. Iwata, M. and T.Hirano (1991) 5. Osmoregulation. in "Fish Physiology", eds. Y.Itazawa and I.Hanyu, Koseisyu Koseikaku, Tokyo, 125-150.
92. Numaguchi, K. (1990) Concentrations of chlorophyll a and phaeo-pigments of bottom sea water, sediment and surface mud in estuary where short necked clam *Ruditapes philippinarum* (Adams et Reeve) inhabited. Bull. Natl. Res. Inst. Aquaculture, 18, 39-50.
93. Numaguchi, K. and J.A.Nell (1991) Effects of gelatin-acacia microcapsule and algal meal supplementation of algal diets on growth rates of Sydney rock oyster, *Saccostrea commercialis* (Iredale & Roughley), larvae. Aquaculture, 94, 65-78.

Nansei National Fisheries Research Institute

94. Hamaguchi, M. (1991) Studies on cellular immune response of yellowtail *Seriola quinqueradiata* against pseudotuberculosis. Bull. Nansei Natl. Fish. Res. Inst., 24, 27-151.
95. Hamaguchi, M., H. Usuki and R. Kusuda (1991) Pasteurella piscicida infection in Yatabe blenny, *Pictiblennius yatabei*. Gyogyo Kenkyu, 26(2), 93-94.
96. Edahiro, T., M. Hamaguchi and R. Kusuda (1991) Suppressive effect of glycyrrhizin against streptococcal infection promoted by feeding oxidized lipids to yellowtail *Seriola quinqueradiata*. SUISANZOSYOKU, 39(1), 21-27.
97. Mustahal and T. Honjo (1988) Growth rate and division periodicity of three diatom species collected from Gokasho Bay, Japan. J. Penel. Budidaya Pantai, 4(2), 46-52.
98. Imai, I., S. Itakura and K. Itoh (1990) Distribution of diatom resting cells in sediments of Harima-Nada and northern Hiroshima Bay, the Seto Inland Sea, Japan. Bull. Coastal Oceanography, 28(1), 75-84.
99. Imai, I., S. Itakura and K. Itoh (1991) Life cycle strategies of the red tide causing flagellates *Chattonella* (Raphidophyceae) in the Seto Inland Sea. Marine Pollution Bulletin, 23, 165-170.
100. Imai, I. and S. Itakura (1991) Densities of dormant cells of the red tide flagellate *Heterosigma akashiwo* (Raphidophyceae) in bottom sediments of northern Hiroshima Bay, Japan. Bull. Japan. Soc. Microbial Ecol., 6(1), 1-7.
101. Imai, I., Y. Ishida, S. Sawayama and Y. Hata (1991) Isolation of a marine gliding bacterium that kills *Chattonella antiqua* (Raphidophyceae). Nippon Suisan Gakkaishi, 57(7), 1409.
102. Itakura, S., I. Imai and K. Itoh (1991) Spatial distribution of cysts of the noxious red tide flagellates *Chattonella* (Raphidophyceae) in sediments of the Eastern Seto Inland Sea, Japan. Nippon Suisan Gakkaishi, 57(6), 1079-1088.
103. Tamai, K. (1990) Tolerance of benthic animals to oxygen deficiency. in "Coastal Fish Farms and the Environment", ed. T. Watanabe, Koseisha Koseikaku, Tokyo, 69-78.
104. Nagai, T. (1991) Trends on catch and stock abundance indices of the Japanese anchovy in the Seto Inland Sea. Bull. Nansei

Natl. Fish. Res. Inst., 24, 1-26.

105. Ogawa, Y., H. Hashimoto, S. Kakuda and K. Gushima (1991) On the growth and life span of the population of oriental river prawn *Macrobrachium nipponense* (De Haan) in the Ashida river. J. Fac. Appl. Biol. Sci. Hiroshima Univ., 30, 43-53.
106. Sato, R., T. Tokai and H. Sakaji (1991) On the existence of a few subpopulations of tiger puffer *Takifugu rubripes* in the Seto Inland Sea. Gyogyo Shigen Kenkyukaigi Nishi-nippon Sokouo Bukaihou, 19, 77-89.
107. Uchida, T. (1991) Sexual reproduction of *Scrippsiella trochoidea* isolated from Muroran harbor, Hokkaido. Nippon Suisan Gakkaishi, 57(6), 1215.
108. Yamaguchi, M. and T. Honjo (1990) Phased cell division in the noxious red tide flagellate *Gymnodinium nagasakiense* (Dinophyceae) and application of the frequency of dividing cells technique for the determination of growth rate. Bull. Plankton Soc. Japan, 37(1), 3-18.
109. Yamaguchi, M., I. Imai and T. Honjo (1991) Effects of temperature salinity and irradiance on the growth rates of the noxious red tide flagellates *Chattonella antiqua* and *C. marina* (Raphidophyceae). Nippon Suisan Gakkaishi, 57(7), 1277-1284.

Seikai National Fisheries Research Institute

110. Okauchi, M., W-J. Zhou, W-H. Zou, K. Fukusho and A. Kanazawa (1990) Difference in nutritive value of a microalga *Nannochloropsis oculata* at various growth phases. Nippon Suisan Gakkaishi, 56(8), 1293-1298.
111. Okauchi, M. (1990) Food value of *Isochrysis* aff. *galbana* for the growth of pearl oyster spat. Nippon Suisan Gakkaishi, 56(8), 1343.
112. Yamasaki, M. (1991) Food consumption of juvenile halibutids in laboratory experiments. Nippon Suisan Gakkaishi, 57(5), 865-867.

Research Division, Fisheries Agency

113. Annual Report on Japan's Fisheries, Fiscal 1990. The Ministry of Agriculture, Forestry and Fisheries, Government of Japan.